

Références

1. Pippias M, Jager KJ, Kramer A, Leivestad T, Sánchez MB, Caskey FJ, et al. The changing trends and outcomes in renal replacement therapy: data from the ERA-EDTA Registry. *Nephrol Dial Transplant*. 2016;31(5):831-41.
2. MacPhail A, Ibrahim JE, Fetherstonhaugh D, Levidiotis V. The Overuse, Underuse, and Misuse of Dialysis in ESKD Patients with Dementia. *Semin Dial*. oct 2015;28(5):490-6.
3. Treatment decisions for older adults with advanced chronic kidney disease | BMC Nephrology | Full Text [Internet]. [cité 3 oct 2019]. Disponible sur: <https://bmcnephrol.biomedcentral.com/articles/10.1186/s12882-017-0617-3>
4. Farrington K, Covic A, Nistor I, Aucella F, Clyne N, De Vos L, et al. Clinical Practice Guideline on management of older patients with chronic kidney disease stage 3b or higher (eGFR<45 mL/min/1.73 m²): a summary document from the European Renal Best Practice Group. *Nephrol Dial Transplant*. 1 janv 2017;32(1):9-16.
5. Carson RC, Juszczak M, Davenport A, Burns A. Is maximum conservative management an equivalent treatment option to dialysis for elderly patients with significant comorbid disease? *Clin J Am Soc Nephrol*. oct 2009;4(10):1611-9.